

Case Report **Published Date:-2018-06-26 00:00:00**

[Low vision due to persistent Cloquet's canal](#)

The Cloquet's canal (CC) is the remnant of the primary vitreous and it disappears in the intrauterine developmental period of an eye. Rarely it can persist in mature eyes and cause low visual acuity. We detected a persistent CC and low visual acuity in a 36 year-old patient. Fundus fluorescein angiography and optical coherence tomography examinations were completely normal in our case and low visual acuity was directly associated with opacification of the fibrovascular persistent CC tissue. Persistent CC can cause low visual acuity due to directly media opacity effect of fibrovascular tissue without any complication.

Review Article **Published Date:-2018-03-23 00:00:00**

[Treatment of Retinal Pigment Epithelial Detachment](#)

Anatomical separation of the retinal pigment epithelium from the Bruch membrane is defined as retinal pigment epithelial detachment (PED) and it is classified as drusenoid, serous, and vascularized. Vascularized PED is mostly associated with choroidal neovascular membrane due to age-related macular degeneration and the risk of vision loss is high in this situation. Studies show that all of baseline values including BCVA, PED height, subretinal fluid, central macular thickness, PED volume, vertical dimension, presence of coincident macular pathology, reflectivity and morphology on optical coherence tomography have prognostic importance. Current treatment protocols mainly based on intravitreal injection of anti-vascular endothelial growth factor (VEGF). Even the bevacizumab was the first anti-VEGF that was used for treatment in PED, there are several reports show the insufficiency of bevacizumab. In treatment-naïve eyes, both of ranibizumab and aflibercept have similar effect in vascularized PED. In treatment-resistant eyes, high dose bevacizumab or switching therapy of anti-VEGF procedures can be effective when considering of all cases, aflibercept seems more effective than other options. We aimed in this manuscript, to give a general information about different characteristics of PEDs and to investigate the treatment strategies in the light of current literature.

Opinion **Published Date:-2018-02-09 00:00:00**

[Place of beta-radiation in the etiology and treatment of cataract](#)

Among eye diseases, cataract is the most commonly encountered lens disease and the leading cause of reduced vision. Cataract caused by radiation develops due to neck & head, central nervous system tumors, eye localized tumors and total body irradiation. Today, the only treatment of cataract is surgery.

Beta radiation is seen to have an important place both in the etiology and treatment of cataract. Beta-radiation creates cataract in the lens as an adverse effect. However, beta radiation implementation is used for delay or prevention of cataract in glaucoma surgery. Effects of beta-radiation on the etiology and treatment should be supported by further prospective clinical studies.

Case Report **Published Date:-2018-02-07 00:00:00**

[The management of Irvine-Gass Syndrome in a patient using Inhaler Steroid](#)

Irvine-Gass syndrome, is one of the most common causes of painless decrease in vision following even uneventful cataract surgery. It usually responds well to medical therapy, but, there are no widely accepted consensus on the efficacy of various therapeutic options for the treatment of Irvine-Gass syndrome. The patient presenting in this case report, has systemic hypertension and chronic obstructive pulmonary disease and he use oral anti-hypertension medication and inhaler steroid. He diagnosed as Irvine-Gass syndrome due to presence of decrease in visual acuity and macular edema with hyporeflexive cystic intraretinal spaces in optical coherence tomography (OCT) since 4th week control visit following uneventful cataract surgery. After the responsiveness of several medications including topical steroid and non-steroidal anti-inflammatory drugs and intravitreal injection of anti-vascular endothelial growth factor (anti-VEGF), intravitreal sustained-release dexamethasone implant was applied. The visual acuity improved to 0.00 logMAR at 1st month after intravitreal dexamethasone therapy and consecutive OCT images showed complete resolution of macular edema with a normalization of the foveal profile. The visual acuity and foveal architecture remained stable in 2-year follow-up period and additional treatment was not needed. To the best of our knowledge, this is the first report that mentions the increment of visual acuity after a single dexamethasone implant, even though it did not response anti-VEGF combined with topical steroid and non-steroidal anti-inflammatory drugs.
